



# Gut microbiome composition better reflects host phylogeny than diet diversity in breeding wood-warblers

Marcella D. Baiz<sup>1</sup>  | Andrea Benavides C.<sup>1</sup> | Eliot T. Miller<sup>2</sup> | Andrew W. Wood<sup>1</sup> | David P. L. Toews<sup>1</sup> 

<sup>1</sup>Department of Biology, Pennsylvania State University, University Park, Pennsylvania, USA

<sup>2</sup>Cornell Lab of Ornithology, Ithaca, New York, USA

## Correspondence

Marcella D. Baiz, Department of Biology, Pennsylvania State University, University Park, PA, USA.

Email: [marcelladbaiz@gmail.com](mailto:marcelladbaiz@gmail.com)

## Funding information

Cornell Lab of Ornithology; National Science Foundation, Grant/Award Number: 2010679; Pennsylvania State University

Handling Editor: Camille Bonneaud

## Abstract

Understanding the factors that shape microbiomes can provide insight into the importance of host–symbiont interactions and on co-evolutionary dynamics. Unlike for mammals, previous studies have found little or no support for an influence of host evolutionary history on avian gut microbiome diversity and instead have suggested a greater influence of the environment or diet due to fast gut turnover. Because effects of different factors may be conflated by captivity and sampling design, examining natural variation using large sample sizes is important. Our goal was to overcome these limitations by sampling wild birds to compare environmental, dietary and evolutionary influences on gut microbiome structure. We performed faecal metabarcoding to characterize both the gut microbiome and diet of 15 wood-warbler species across a 4-year period and from two geographical localities. We find host taxonomy generally explained ~10% of the variation between individuals, which is ~6-fold more variation of any other factor considered, including diet diversity. Further, gut microbiome similarity was more congruent with the host phylogeny than with host diet similarity and we found little association between diet diversity and microbiome diversity. Together, our results suggest evolutionary history is the strongest predictor of gut microbiome differentiation among wood-warblers. Although the phylogenetic signal of the warbler gut microbiome is not very strong, our data suggest that a stronger influence of diet (as measured by diet diversity) does not account for this pattern. The mechanism underlying this phylogenetic signal is not clear, but we argue host traits may filter colonization and maintenance of microbes.

## KEYWORDS

16S, birds, COI, metabarcoding, Parulidae, phylosymbiosis

## 1 | INTRODUCTION

Microorganisms that form intimate associations with their hosts can take part in important physiological functions. In particular, the gut microbiome—the community of microbes that colonize the gastrointestinal tract—has been linked to host behaviour, immune function, development, metabolism and disease (Bodawatta, Hird, et al., 2021; Sommer & Bäckhed, 2013; Suzuki, 2017).

The taxonomic composition of the gut microbiome can vary, sometimes dramatically within and between host species (Grond et al., 2019; Loo et al., 2019; Song et al., 2020), as well as within individuals over short timescales (Skeen et al., 2021; Videvall et al., 2019). However, when host–microbe associations are long-term, gut microbiomes may be expected to be species-specific and their assembly to be dependent on host evolutionary divergence (Brooks et al., 2016). Consistent with this, host evolutionary history, in addition to diet,

has been implicated as one of the strongest factors driving vertebrate gut microbiome similarity (Youngblut et al., 2019).

Recent studies have strongly supported a positive correlation between host species divergence and gut microbiome divergence—known as “phylosymbiosis”—particularly for insects and nonflying mammals (Brooks et al., 2016; Song et al., 2020). However, in birds, differences in gut microbiome structure between species are less pronounced (Song et al., 2020). Despite species-level differences in the gut microbiota of 37 New Guinean passerine species (14 families), Bodawatta, Koane, et al. (2021) did not find an influence of host phylogeny on gut microbiome structure. This is in contrast, however, to a study on 51 passerine species (21 families) breeding in the Czech Republic (Kropáčková et al., 2017) and a study on all 15 crane species (family Gruidae) in captivity which found a weak influence of host phylogeny, and only when examining female individuals (Trevelline et al., 2020).

A favoured hypothesis to explain this marked difference in phylosymbiosis between bird and nonflying mammal gut microbiota is that because birds evolved a reduced and simplified gastrointestinal tract as an adaptation to flight, they have highly reduced gut retention times from consumption of food to defecation (Song et al., 2020). This reduced retention time and simplified gut environment may favour high turnover in the avian gut microbiome, and a larger role of the diet and environment over host taxonomy in the structuring of the gut microbiome (Bodawatta, Koane, et al., 2021).

In Darwin's finches, gut microbiome communities cluster more strongly by host habitat than by host species (Loo et al., 2019). Both host phylogeny and diet in this group, which is known for adaptive divergence in beak morphology linked to foraging ecology, show a moderate influence on gut microbiome variation (Loo et al., 2019). Further, the gut microbiome of the vampire finch, a diet specialist, is highly divergent from other species (Michel et al., 2018). Similarly, captive birds tend to have distinct gut microbiota from their wild counterparts, probably resulting from artificial diets, built enclosures and human interaction (San Juan et al., 2021). These studies support a strong role of the environment, including diet, in shaping the avian gut microbiome.

Although many studies have detected effects of diet on the avian gut microbiome (Bodawatta, Koane, et al., 2021; Davidson et al., 2020; Knutie, 2020; Teyssier et al., 2020; Xiao et al., 2021), few have analysed host diet beyond broad categorizations of diet type (e.g., omnivore vs. insectivore) and/or included birds that were fed standardized and non-natural diets (but see Bodawatta et al., 2022; Schmiedová et al., 2022). Further, many studies that have assessed species-specific differences in gut microbiome structure have had limited sample sizes including only one or a few individuals per species or included data collected and sequenced at different times or in different ways. To gain a holistic picture of the effects of host diet, evolutionary history and geography on gut microbiome structure, it will be necessary to sample natural populations using standardized methods. Understanding the factors that shape the avian gut microbiome is important for understanding host-symbiont interactions and co-evolutionary dynamics, and how these dynamics may differ from other taxonomic groups of animals


(i.e., mammals). The role of the gut microbiome in host evolutionary processes is largely unexplored and its potential role in facilitating and responding to avian host adaptive radiation—where species diversification is tied to ecological differentiation—is a major outstanding question (Bodawatta, Hird, et al., 2021).

Here, we characterize the gut microbiome of wood-warblers (family: Parulidae) breeding in sympatry in Eastern North America across a 4-year period and examine factors that may play a role in shaping gut microbiome structure. Parulidae is a passerine radiation of >100 insectivorous species that evolved rapidly in the last 7 million years (Barker et al., 2015; Lovette et al., 2010), and is a classic model for studies of ecological differentiation, including diet niche partitioning (MacArthur, 1958). In the current study, we use 16S faecal metabarcoding to examine gut microbiomes of 15 species representing seven genera (Figure 1a). Our aims are to characterize the “core” parulid gut microbiota (a common set of microbes across individuals) and to quantify differences in gut microbiome composition between hosts. We predict that due to genetic and ecological differentiation among host species, variation in the gut microbiome will be largely explained by host taxonomy. Further, we explicitly test the prediction of phylosymbiosis, where host phylogenetic relatedness should correlate with gut microbiome similarity. We also examine the relationship between gut microbiome diversity and diet diversity by analysing COI metabarcoding sequences amplified from faecal samples of these same individuals. With the presumption that a diet characterized by a high diversity of arthropods will incur ingestion of a greater diversity of bacteria—either associated with arthropod hosts, or the environments in which they are found—we predict that diversity of the warbler gut microbiome and diet will be positively correlated. Finally, we test for other environmental signals in the structuring of host gut microbiomes by examining effects of sampling year, locality and diet specialization.

## 2 | MATERIALS AND METHODS

### 2.1 | Sample collection and DNA extraction

We used mist-nets to capture birds during four consecutive breeding seasons (May–July 2017–2020). In all years, we targeted sampling locations in northern hardwood forests, both in Adirondack Park, New York, and in 2019 and 2020, we also sampled birds in central Pennsylvania (Figure S1; Table 1). We selected sites where a diversity of warbler species (up to eight) could be heard singing so as to maximize sympatry among species included in the study. Upon capture, we held individuals inside a brown paper bag for up to 10 min to allow ample time for excretion inside the bag before removal and subsequent banding. We removed faeces by scraping it from the inside of the bag directly into a sample tube containing lysis buffer (100 mM Tris pH 8; 100 mM Na<sub>2</sub>EDTA, 10 mM NaCl; 0.5% sodium dodecyl sulphate; White & Densmore, 1992), and froze samples at –20°C within 2 weeks of collection. Because we were interested in variation among individuals, we chose a single sample at random to



**FIGURE 1** (a) Phylogenetic relationships between host species in this study. Upside down triangles indicate low-diversity diet, triangles indicate high-diversity diet, and squares indicate intermediate-diet diversity based on our COI diet index. The full data set represents all samples collected between 2017 and 2020, and batch 1 represents all samples collected between 2017 and 2019. Illustrations © Lynx Edicions. (b) Relative abundance of bacterial phyla in the full 16S data set, and (c) relative abundance of arthropod orders in the full COI data set. See Table 1 for host species codes. [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/mec.16762)]

include in our analyses from individuals that were recaptured in the same or subsequent years. In total, we sequenced samples from 408 individuals.

We extracted total DNA from faecal samples using an SPRI-bead faecal DNA extraction method modified from Vo and Jedlicka (2014). Samples were processed in two sets: those collected in 2017–2019 and in 2020. After thawing faecal samples at room temperature, we centrifuged sample tubes and used bleach-sterilized laboratory spatulas after being thoroughly dried and/or pipetting to transfer ~5 mg of faecal material into 2-ml screw-cap microcentrifuge tubes each containing 0.25 g of 0.1-mm and 0.25 g of 0.5-mm zirconia-silica beads. For samples that amounted to <5 mg of faecal material, we supplemented with a suitable volume of storage buffer from inside the sample tube as necessary. We immediately added 818 µl of warmed (65°C) lysis buffer (Vo & Jedlicka, 2014) and homogenized samples using a Precellys 24 Tissue Homogenizer (Bertin Instruments) set to three cycles of 6800 rpm for 30 s with a 30-s pause between cycles. After transferring the supernatant to clean microfuge tubes, we incubated samples with Qiagen Solution C3 (Qiagen DNeasy PowerSoil 12888-100-3) to

remove PCR (polymerase chain reaction) inhibitors. Next, we removed DNA from the supernatant using homemade solid phase reversible immobilization (SPRI) magnetic beads (“Serapure” beads; Rohland & Reich, 2012). Serapure beads were added at a 1.9× bead-to-supernatant volume ratio and, after cleaning with 80% ethanol, we eluted DNA in 10 mM Tris-HCl. Extracted DNA was stored at –20°C before proceeding with library preparation.

We sequenced two types of negative controls. First, negative extraction controls followed the same extraction procedure described above for which the input was sample storage buffer taken from tubes that were transported to the field, but were not used for collecting faecal material. Second, we included PCR-negative controls, which are described below.

## 2.2 | 16S and COI amplicon sequencing

As with DNA extractions, we prepared and sequenced metabarcoding libraries in two separate batches: (i) samples collected between 2017 and 2019, and (ii) samples collected in 2020.

We used a two-step multiplex dual-index amplicon approach to separately prepare 16S libraries and COI libraries for sequencing again following Vo and Jedlicka (2014) with some adjustments. We first used universal 515F/806R primers to amplify the V4 region of the bacterial 16S rRNA gene (Caporaso et al., 2012) and the "ANML" general arthropod COI mitochondrial primers LCOI-1490/COI-CFMRa described in Jusino et al. (2019). Each primer pair was modified with overhanging Illumina adapter sequences. Prior to PCR, we randomized the order of samples to be amplified to avoid within-plate batch effects during amplification. Negative PCR controls were included on each plate. In addition to our faecal samples, we sequenced four negative controls per primer pair in each library pool, with the exception of the first batch COI library pool which did not contain any negative controls. Negatives included two "extraction controls" amplified and sequenced from DNA extractions made from sample tubes containing only buffer (and no faeces) as well as two negative PCR controls.

We performed initial 16S PCR amplification for each sample in triplicate in 30- $\mu$ l reactions comprising 0.2  $\mu$ l Platinum II *Taq* Hot Start DNA Polymerase (Invitrogen 14966005), 5  $\mu$ l 5 $\times$  Buffer (Invitrogen 14966005), 1.25  $\mu$ l of each primer (10  $\mu$ M concentration), 13.5  $\mu$ l molecular-grade water, 0.5  $\mu$ l 10 mM dNTP mix (Promega U151A) and 3.3  $\mu$ l of faecal DNA. Reaction conditions followed the two-step PCR protocol recommended by the manufacturer: 94°C for 2 min, followed by 34 cycles of 98°C for 5 s, 68°C for 15 s, followed by a final extension at 68°C for 5 min, and hold at 12°C. We performed initial COI PCR amplification in 30- $\mu$ l reactions comprising 0.24  $\mu$ l Platinum II *Taq* Hot Start DNA Polymerase, 6  $\mu$ l 5 $\times$  Buffer, 1.5  $\mu$ l of each primer (10  $\mu$ M concentration), 16.16  $\mu$ l molecular-grade water, 0.6  $\mu$ l 10 mM dNTP mix and 4  $\mu$ l of faecal DNA. Reaction conditions followed Jusino et al. (2019) with minor adjustments: 94°C for 2 min, followed by five cycles of 94°C for 15 s, 45°C for 15 s, 68°C for 15 s, followed by 35 cycles of 98°C for 5 s, 68°C for 15 s, followed by a final extension at 68°C for 5 min, and hold at 12°C. We cleaned initial PCR products by incubating with a 1 $\times$  volume of Serapure beads and eluting the bound DNA in 10 mM Tris-HCl. Triplicate 16S reactions were pooled before this cleaning step. Then we evaluated amplification success by visualizing the cleaned product on a 1.5% agarose gel.

Next, we appended dual P5 and P7 Illumina indexes to each library via PCR. Reactions were 30  $\mu$ l and contained 15  $\mu$ l KAPA HiFi HotStart ReadyMix (Roche 7958935001), 3  $\mu$ l of each primer (10  $\mu$ M concentration) and 9  $\mu$ l DNA (cleaned initial PCR product). Reaction conditions followed the manufacturer's recommendations: 98°C for 45 s, followed by five cycles of 98°C for 15 s, 60°C for 15 s, 72°C for 15 s, followed by a final extension at 72°C for 1 min and hold at 12°C. We then cleaned the indexed PCR product using a double-sided Serapure bead procedure. We first removed potential high-molecular-weight contamination by incubating PCR product with a 0.75 $\times$  volume of Serapure beads. After placing the samples on the magnet, we transferred the supernatant to fresh tubes and incubated it with a 1 $\times$  volume of Serapure beads to remove potential low-molecular-weight contamination. DNA was

eluted in 10 mM Tris-HCl, and we evaluated amplification success as for the initial PCR.

We quantified DNA in our final PCR products with a Qubit 4.0 Fluorometer (Invitrogen). We then normalized library concentrations and pooled libraries to a final pool concentration of at least 2 nM. We submitted the final pool to the Penn State Genomics Core Facility to perform final quality assessment on a Bioanalyzer Tape Station and confirm pool concentration with quantitative (q) PCR. Samples were then sequenced with an Illumina MiSeq using the 600-cycle kit run as 250 $\times$ 250 paired-end sequencing.

For the first batch of samples, 16S and COI libraries were independently pooled and each pool was sequenced in a single lane of Illumina sequencing. The second batch included a smaller number of samples, so to achieve a similar depth of sequencing as in the first batch, we pooled and sequenced 16S libraries and COI libraries together in the same sequencing lane.

### 2.3 | 16S amplicon sequence processing

We used QIIME 2 version 2020.8 (Bolyen et al., 2019) to process 16S sequencing reads and obtain a table of counts of amplicon sequence variants (ASVs, or amplicon sequences representing microbial taxonomic units) across samples. For each sequencing run, we imported demultiplexed paired-end sequences, used the function *qiime dada2 denoise-paired* to trim primer sequences from the 3' ends of reads, and to trim five bases from the 5' ends of reads before merging read pairs and detecting ASVs. We then assigned taxonomic classification to ASVs using the SILVA database (version 138 SSURef NR99, Quast et al., 2012).

Upon classification, we removed mitochondrial, chloroplast, unassigned and eukaryotic ASVs. We also identified and removed possible contaminant ASVs by contrasting the presence/absence of ASVs in our negative controls with their prevalence in experimental faecal samples (i.e., non-negative controls) using the R package *decontam* (Davis et al., 2018). We used the "prevalence" method to identify and remove ASVs more prevalent in negative controls than in experimental samples using a probability threshold of 0.5. We also manually removed ASVs present in negative controls, but absent in experimental samples, as these were also probably contaminants. In total, we removed 87 and 359 contaminant ASVs from the batch 1 and batch 2 data sets, respectively.

At this point, we used QIIME 2 to merge the feature table, representative sequences and taxonomy files from the two separate sequencing runs. We finally generated a phylogenetic tree from the merged set of ASV sequences for downstream diversity analyses. We used *qiime phylogeny align-to-tree-mafft-fasttree* to perform multiple sequence alignment, mask highly variable positions, and first generate an unrooted tree and finally a tree rooted at the midpoint of the longest tip-to-tip distance of the unrooted tree.

Finally, we applied several additional filtering steps to achieve a high-quality representation of warbler gut microbiomes. We excluded individuals from species represented by fewer than five individuals

TABLE 1 Sampling information for species included in this study

Species code	English common name	Latin name	2017		2018		2019		2020		Total	
			NY	PA	NY	PA	NY	PA	NY	PA	Subset batch 1	Full data set
AMRE	American redstart	<i>Setophaga ruticilla</i>	2	0	3	0	5	6	4	4	16	23
BAWW	Black-and-white warbler	<i>Mniotilta varia</i>	0	0	7	0	1	1	4	9	9	22
BLBW	Blackburnian warbler	<i>Setophaga fusca</i>	1	0	4	0	3	1	5	6	9	20
BTBW	Black-throated blue warbler	<i>Setophaga caerulescens</i>	2	0	4	0	2	5	5	10	13	28
BTNW	Black-throated green warbler	<i>Setophaga virens</i>	3	0	4	0	4	3	5	7	14	26
CAWA	Canada warbler	<i>Cardellina canadensis</i>	2	0	4	0	3	3	4	1	12	17
COVE	Common yellowthroat	<i>Geothlypis trichas</i>	1	0	4	0	3	1	3	2	9	14
CSWA	Chestnut-sided warbler	<i>Setophaga pensylvanica</i>	5	0	2	0	1	5	3	3	13	18
HOWA	Hooded warbler	<i>Setophaga citrina</i>	0	0	0	0	0	6	0	5	6	11
MAWA	Magnolia warbler	<i>Setophaga magna</i>	1	0	6	0	4	1	4	1	12	17
MYWA	Myrtle warbler	<i>Setophaga coronata coronata</i>	4	0	5	0	5	1	5	0	15	20
NAWA	Nashville warbler	<i>Leiothlypis ruficapilla</i>	2	0	4	0	5	0	0	0	11	11
NOPA	Northern parula	<i>Setophaga americana</i>	2	0	4	0	5	0	5	0	11	16
OVEN	Ovenbird	<i>Seiurus aurocapilla</i>	2	0	4	0	4	1	5	6	11	22
WEWA	Worm-eating warbler	<i>Helminthos vermivorum</i>	0	0	0	0	0	0	0	5	0	5
Total individuals											161	270

Note: Subset batch 1 includes samples collected between 2017 and 2019, which were sequenced together in a single sequencing run.

Abbreviations: NY, New York; PA, Pennsylvania.

in our data set because we were interested in examining species differences in gut microbiome structure. Because very low depth and uneven depth of sequencing among samples can affect diversity estimates (Hughes & Hellmann, 2005), we next generated a rarefied data set by randomly downsampling ASVs to a minimum threshold to standardize total read counts across samples. We determined the minimum acceptable ASV count threshold by examining rarefaction curves constructed using the *rarecurve* function in *vegan* (Oksanen et al., 2020) using a step size of 50. Based on this analysis, we determined a library size of 4000 reads to be an acceptable threshold since the number of observed ASVs appears to plateau beyond this point (Figure S2a).

Because we detected a significant effect of sequencing batch on our diversity estimates (i.e., a “batch effect”; see Results), we also performed analyses on a subset of the data that only included the first batch of samples (collected between 2017 and 2019, referred to as “batch 1”). For these analyses, we performed the same sequence processing steps as above except for merging-in data from the samples collected in 2020.

## 2.4 | COI amplicon sequence processing

We used the *AMPTK* (version 1.5.3) pipeline to analyse COI metabarcoding data by applying the default clustering algorithm (*VESEARCH* version 2.17.1) for operational taxonomic units (OTUs) and assigned taxonomy by pulling from the chordates and arthropods in the BOLDv4 database. We rooted the OTU phylogeny output from *AMPTK* on a randomly chosen arachnid OTU, as arachnids split from the common arthropod ancestor prior to insects. We then imported the COI metabarcoding data into *PHYLOSEQ* for downstream analyses and applied a similar framework as we did with our 16S data. We first removed OTUs assigned to the phylum Chordata as this represents off-target amplification, then rarefied depth to 15,000 reads per individual (full data set), and 8500 reads per individual (batch 1 subset; Figure S2b).

For analyses where we directly investigated the effect of diet on the microbiome at the individual level, we only analysed individuals with data that passed filtering steps in both microbiome and diet data sets. This included 216 individuals in the full data set representing 15 species (mean = 14 individuals per species, *SD* = 4.7) and 130 individuals in the batch 1 subset representing 14 species (mean = 9 individuals per species, *SD* = 1.7).

## 2.5 | Diet diversity and its relationship with gut microbiome diversity

We estimated within-individual diversity (alpha diversity) of the diet and gut microbiome using the Shannon index and the Chao1 index using the *diversity* function in *vegan*, and using Faith's phylogenetic diversity using the *estimate\_pd* function in *btools* (Battaglia, 2022). The Shannon index quantifies ASV richness (the number of ASVs) as well as evenness (the equity in ASV abundances), while Chao1 just quantifies ASV richness. Faith's phylogenetic diversity is a measure

of ASV richness that is the sum of branch lengths in the phylogeny that connect all ASVs in the community assemblage. We estimated between-individual differences between microbiomes (beta diversity) using four different metrics: Bray–Curtis, Jaccard, unweighted UniFrac and weighted UniFrac, calculated using the *distance* function in *phyloseq* (McMurdie & Holmes, 2013). Bray–Curtis measures differences in community composition and is based on ASV abundances, whereas Jaccard is based only on presence/absence and does not rely on abundance. Unweighted UniFrac measures the phylogenetic distance between communities based on presence/absence of ASVs, whereas weighted UniFrac is similar but weights branch lengths by ASV abundance.

We used three approaches to examine the relationship between diet and the gut microbiome. With the prediction that a generalized diet, characterized by a high diversity of arthropod taxa, supports a high gut microbiome diversity, we first tested for a positive correlation between individual diet alpha diversity and gut microbiome alpha diversity using a Kendall's rank correlation test. To account for phylogenetic history, we also ran phylogenetic linear models to test for a linear relationship between mean within-species alpha diversities of the gut microbiome and diet using *phylolm* (Tung Ho & Ané, 2014). In each model, we included the host phylogeny with branch lengths scaled using divergence times (see below) and the lambda model of covariance, using 1000 bootstrap replicates.

Second, at the species level, we tested whether gut microbiome structure differs among species with a more specialized and less diverse diet, and species with a more generalized and more diverse diet using permutational multivariate analysis of variance (PERMANOVA) of beta diversity distances using the *adonis2* function in *vegan*. For this analysis, we categorized each species as either a “low diversity” diet, “high diversity” diet or “intermediate” by creating an index of diet specialization (Figure 1a). To calculate this index, we summed mean individual within-species diet alpha diversity and mean within-species diet beta diversity with the assumption that (i) more specialized diets are characterized by a lower diversity of food items (low alpha diversity) and individuals within more specialized species eat a similar diet (low beta diversity), and (ii) more generalized diets are characterized by a high diversity of food items (high alpha diversity), and individuals within more generalized species may have highly divergent diets depending on local food availability (high beta diversity). Thus, a low score reflects a less diverse and more specialized diet, and a high score reflects a more diverse and more generalized diet. We note that this index quantifies diversity of the diet and that host species within the same diet categorization may have dissimilar diets by way of diet content (e.g., proportion that is flying insects).

For both alpha diversity and beta diversity of the diet, the different diversity metrics we calculated were positively correlated (with the exception of weighted UniFrac and unweighted UniFrac beta distance when using the full data set; Table S1) and diet type classification of each species was consistent across metrics. Thus, for simplicity we report the index of diet specialization using the

Shannon index to estimate alpha diversity and the Bray–Curtis metric to estimate beta diversity:

$$\text{index of diet specialization} = \text{mean diet } \alpha_{\text{Shannon}} + \text{mean diet } \beta_{\text{Bray-Curtis}}$$

We note that because we used a subset of individuals to calculate diet index for batch 1, for some species classification of diet diversity using the diet index is not consistent between this subset and the full data set. Four species are classified as intermediate in one data set and either as high diversity or as low diversity in the other data set. However, species diet index values are positively correlated between the full data set and batch 1 ( $\tau = 0.516$ ,  $p = .010$ , Figure S3), suggesting this index is robust to individual variation in diet. Our results do not change when excluding these four species from the analyses so we include them in our analyses.

Finally, we used topological congruence analysis to determine whether similarity in gut microbiome structure among host species reflects diet similarity with the expectation that if diet directly shapes host gut microbiomes, then clustering of species by diet similarity will mirror clustering of species by gut microbiome similarity. To generate dendrograms representative of each species, we generated a new ASV table—each for rarefied COI sequence counts and rarefied 16S sequence counts—grouped by host species by averaging ASV counts within each species, recalculated dissimilarity matrices and constructed dendrograms by clustering distance matrices using the UPGMA method in the *hclust* function in R (following Trevelline et al., 2020). We then compared the observed 16S dendrogram to the observed COI dendrogram using *TreeCmp* (Bogdanowicz et al., 2012) to compute the matching cluster metric of topological congruence (Bogdanowicz & Giaro, 2013). Following Brooks et al. (2016), we then compared the observed 16S dendrogram with 10,000 dendrograms with randomized topology and calculated a normalized congruence score, which is the observed matching cluster score divided by the maximum congruence score between the observed dendrogram and one of the random dendrograms. Finally, we evaluated significance and report a  $p$ -value by dividing the number of randomized dendrograms with equal or greater congruent scores to the observed 16S dendrogram than the score between the two observed dendrograms by 10,000. We also used Mantel tests as a complementary analysis to examine correlations between the diet and microbiome beta distance matrices at the individual level, where each value represents the beta distance between a pair of individuals, using *vegan::mantel* with the Spearman correlation method.

## 2.6 | Gut microbiome diversity and topological analyses

We identified a “core” wood-warbler gut microbiome as the collection of ASVs present across a large number of individuals using the rarefied data set. Because most ASVs had a low prevalence among individuals (Figure S4), we report the core microbiome as ASVs present in >30% of all individuals. Although this threshold is arbitrary, we believe it is conservative as only 39 ASVs were represented in more than 30% of individuals (see below). We also report taxa at

high relative abundance across all samples at the phylum level. This set of ASVs represents bacteria that are most common in the gut microbiome among breeding male wood-warblers.

To quantify the effect of host taxonomy on the gut microbiome and the extent to which gut microbiomes covary with host phylogeny, we took two approaches using the full set of ASVs. First, we estimated gut microbiome divergence (beta diversity) among individuals using four measures of community dissimilarity: Bray–Curtis distance, Jaccard distance, and weighted and unweighted UniFrac distances. We then used *vegan::adonis2* to conduct PERMANOVA tests to determine the effect of host species on community dissimilarity. Because our samples were collected across four breeding seasons, from two geographical localities, and were sequenced in two different batches we also tested for effects of these factors. We included each of these factors in our model and set the “by” parameter to “margin.” However, in the full data set the effects of sampling year and sequencing batch are confounded since all samples collected in 2020 were sequenced in batch 2, so we ran two separate models which included either host species + locality + year, or host species + locality + sequencing run. Results for host species and locality were similar between models, so we report results from the model that included sequencing run for simplicity. We also calculated multivariate homogeneity of group dispersions for significant variables using *vegan::betadisper* and assessed deviations from this expectation using *vegan::permutest* because a homogeneous dispersion among groups is an assumption for PERMANOVA tests. We visualized beta distances between gut microbiota using the principal coordinate analysis (PCoA) method of *phyloseq::ordinate*.

Our second approach was to test for congruence between the host phylogeny and microbiome, as phylosymbiosis predicts that host relatedness and microbiome community similarity should exhibit a positive relationship (Brooks et al., 2016). To do this, we first used the same topological congruence approach as described above, but used the topology from Baiz et al. (2021) for *Setophaga* species, and from Lovette et al. (2010) for outgroup taxa in place of the diet dendrogram (Figure 1a). We then also used Mantel tests to test for correlations between the gut microbiome distance matrix and a matrix of cophenetic distances, representing evolutionary distances, between individuals. We calculated cophenetic distance between species using the *stats::cophenetic* function on a dendrogram representing the host phylogeny in Figure 1a, with branch lengths scaled using divergence times from TimeTree of Life (Kumar et al., 2017; Table S2). Note that an evolutionary distance of zero denotes a pair of individuals from the same species.

Because we found a significant influence of sequencing batch on gut microbiome diversity, we separately performed all analyses on the subset of samples sequenced in the first batch (collected between 2017 and 2019, referred to as “batch 1”) as this batch included a larger subset of samples, which were collected across multiple years, than the second batch, which only included samples collected in 2020. For topology and Mantel analyses, we also subset our data to account for potentially confounding effects of (i) geographical locality by only analysing samples collected in New York between 2017 and 2019 (referred to as “batch 1-NY”)

and (ii) sampling year by only analysing samples collected in 2020 (referred to as "batch 2").

### 3 | RESULTS

#### 3.1 | 16S sequencing output and composition of the warbler gut microbiome

The number of ASVs yielded by our first 16S sequencing run was 6412 (per-individual median = 36, mean = 53,  $SD = 65$ ) while our second 16S sequencing run yielded 10,590 ASVs (per-individual median = 235, mean = 218,  $SD = 73$ ). This discrepancy is probably explained by a higher average depth of sequencing across individuals in the second sequencing run (Figure S5), despite our attempt at normalization. Taxa that were detected in both sequencing runs represented a small proportion of the total number of ASVs across runs (6%), contributing to the gut microbiome differentiation we observed for individuals sampled in 2020 (see below).

After merging our 16S data sets, applying our filtering steps and rarefaction, our full data set consists of 270 individuals representing 15 species (mean 18 individuals per species, with 95% of individuals being male, 1% female and 4% of unknown sex). Among these samples, we detected 12,048 ASVs from 39 bacterial phyla with the top phylum, Proteobacteria, representing 60% of the total reads (Figure 1b). Firmicutes was the next most abundant phylum, representing 13% of the total reads, followed by Actinobacteriota, representing 6.5% of the total reads. The remaining phyla each represented <5% of the total reads. We observed considerable variation in relative abundance of prevalent taxa between individuals of the same species (Figure S6a). Despite low overlap in ASV identity between sequencing runs, the composition and relative abundance of prevalent phyla were very similar across host species when we separately examined samples that were sequenced in different batches (Figure S7).

Most ASVs were present in <10% of individuals, and only 39 ASVs were represented in >30% of individuals (Figure S4). Each of these core ASVs was represented in all but one or two of the host species we analysed (Tables S3 and S4). The most prevalent ASV was a Gammaproteobacteria of the family Yersiniaceae. This ASV was found in all 15 host species and ~60% of samples in both the full data set and the batch 1 subset. Gut microbiome alpha diversity did not differ among host species (Kruskal–Wallis rank sum test: Shannon index: full data set  $d.f. = 14$ ,  $\chi^2 = 14.68$ ,  $p = .400$ ; batch 1  $df = 13$ ,  $\chi^2 = 16.354$ ,  $p = .231$ , Chao1 index: full data set  $df = 14$ ,  $\chi^2 = 13.99$ ,  $p = .451$ ; batch 1  $df = 13$ ,  $\chi^2 = 19.32$ ,  $p = .113$ , Faith's PD: full data set  $df = 14$ ,  $\chi^2 = 14.98$ ,  $p = .380$ ; batch 1  $df = 13$ ,  $\chi^2 = 18.764$ ,  $p = .131$ ).

#### 3.2 | COI sequencing output, diet diversity and its relationship with gut microbiome diversity

Our first COI sequencing run yielded 3235 OTUs, while the second yielded 2668 OTUs. In contrast to the 16S data set, there was

moderate overlap in OTU identity between sequencing runs (37% of OTUs are represented in both batches).


Our analyses revealed 4397 OTUs in the full COI data set, which was reduced to 3227 after filtering and rarefaction. Among warbler species, ~70% or greater relative abundance of diet taxa consisted of insects, particularly in the orders Diptera and Lepidoptera (Figure 1c; Figure S6b). The majority of other diet taxa included Arachnids in the family Araneae. There was a high degree of overlap among species in diet PCoA space (Figure 3b). These results were consistent between analyses that included all individuals and only individuals sequenced in the first batch.

Warbler species fell into three natural partitions along our index of diet specialization, and thus we used these partitions to classify species according to diet type (Figure 2b). We classified two of three warbler species with low diversity diets depending on the data set being analysed (batch 1: American redstart [AMRE], chestnut-sided warbler [CSWA]; full data set: American redstart [AMRE], chestnut-sided warbler [CSWA], worm-eating warbler [WEWA]). We classified two to four species with high diversity diets (batch 1: black-throated green warbler [BTNW], Canada warbler [CAWA]; full data set: black-and-white warbler [BAWW], Canada warbler [CAWA], common yellowthroat [COYE], hooded warbler [HOWA]), and the remainder of species as intermediate (Figure 1a).


When considering within-individual diversity, we found no correlation between diet alpha diversity and microbiome alpha diversity when using Shannon index and Faith's PD, as well as Chao1 when considering the full data set (Kendall's rank correlation, Shannon index batch 1:  $\tau = 0.029$ ,  $p = .619$ ; full data set:  $\tau = 0.005$ ,  $p = .906$ , Faith's PD batch 1:  $\tau = 0.110$ ,  $p = .064$ ; full data set:  $\tau = -0.034$ ,  $p = .459$ , Chao1 full data set:  $\tau = -0.063$ ,  $p = .171$ ), but when considering the batch 1 subset using Chao1, alpha diversity of the diet and microbiome were positively correlated (Figure 2a; Kendall's rank correlation batch 1:  $\tau = 0.124$ ,  $p = .038$ ). This indicates that, for batch 1, individuals that consumed high-richness diets (more OTUs) tended to have more rich gut microbiota (more ASVs), although the correlation is weak. After accounting for evolutionary history, our phylogenetic linear models revealed no relationship between mean species gut microbiome alpha diversity and mean species diet alpha diversity (Shannon index batch 1: coefficient = 0.860,  $p = .089$ ; full data set: coefficient = -0.405,  $p = .273$ , Faith's PD batch 1: coefficient = -0.145,  $p = .789$ ; full data set: coefficient = 0.340,  $p = .624$ , Chao1 batch 1: coefficient = 0.305,  $p = .480$ ; full data set: coefficient = -0.024,  $p = .956$ ).

Alpha diversity of the microbiome was generally lower for individuals of species that were diet specialists, and higher for individuals of species that were diet generalists (Figure 2c), but alpha diversity of the microbiome did not differ significantly by species diet type (Kruskal–Wallis  $df = 2$ : Shannon index batch 1:  $\chi^2 = 2.8$ ,  $p = .242$ ; full data set:  $\chi^2 = 0.014$ ,  $p = .993$ , Chao1 batch 1:  $\chi^2 = 5.4$ ,  $p = .068$ ; full data set:  $\chi^2 = 0.31$ ,  $p = .855$ , Faith's PD batch 1:  $\chi^2 = 4.4$ ,  $p = .110$ ; full data set:  $\chi^2 = 0.13$ ,  $p = .936$ ), even when only comparing low-diversity diets to high-diversity diets (Kruskal–Wallis  $df = 1$ : Shannon index batch 1:  $\chi^2 = 1.934$ ,  $p = .1643$ , full data set:





**FIGURE 2** Relationship between diet diversity and gut microbiome diversity. (a) Within-individual diversity of the gut microbiome is weakly correlated with within-individual diet diversity as measured by the Chao1 index in the batch 1 data set. Dashed line is a linear model fit to the data. Point colour in each panel reflects warbler species (see Table 1 for species codes). (b) Distribution of diet index scores by host species, where a low score is reflective of low diet diversity or diet specialization. Colour indicates assignment to diet type and is consistent with box plot colour in panel (c) and with ellipse colour in panel (d). (c) Microbiome alpha diversity does not differ among diet types, as classified by diet index. (d) Principal coordinate analysis (PCoA) of Bray–Curtis distance between host gut microbiomes sequenced in batch 1. Point shape represents species diet type as defined by our index of diet specialization. Ellipses are drawn at 50% confidence level, and line colour indicates diet type. In each panel, data shown are from sequencing batch 1. [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]



**FIGURE 3** Principal coordinate analysis (PCoA) of Bray–Curtis distance between (a) host gut microbiomes and (b) individual diet. In both panels, data are from samples sequenced in batch 1. Point colour represents host species, and shape represents geographical locality. See Table 1 for species codes. [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]

$\chi^2 = 0.049$ ,  $p = .825$ , Chao1 batch 1:  $\chi^2 = 1.8$ ,  $p = .181$ ; full data set:  $\chi^2 = 0.29$ ,  $p = .592$ , Faith's PD batch 1:  $\chi^2 = 1.3$ ,  $p = .259$ ; full data set:  $\chi^2 = 0.17$ ,  $p = .676$ .

### 3.3 | Factors accounting for warbler gut microbiome structure

When analysing the full data set which included microbiomes sequenced in two different sequencing runs, there was a very clear and strong batch effect where microbiomes sequenced in one run were more similar to each other than to microbiomes sequenced in the other run (Figure S8). Yet, PCoA of gut microbiome dissimilarity matrices revealed a high degree of overlap among hosts of different species and among hosts from different geographical localities (Figure 3a). There was little clustering of microbiomes by diet

type of host species as defined by our index of diet specialization (Figure 2d).

Our PERMANOVA tests (Table 2) revealed that sequencing run explained a relatively high degree of variation in Bray–Curtis distances (13%,  $p = .001$ ), Jaccard distances (7.1%,  $p = .001$ ) and unweighted UniFrac distances (7.9%,  $p = .001$ ). This strong batch effect probably confounded tests of other variables, since the second sequencing run only contained samples collected in a single year (2020) and included an additional species (WEWA, worm-eating warbler) that is not represented in the first sequencing run. Thus, we analysed the subset of data from 2017–2019 (i.e., batch 1) separately to examine the effect of biological factors on microbiome structure in the absence of the sequencing batch effect, because of the two sequencing runs this batch included the largest sample size of individuals and included 3 years of sampling. This analysis revealed sampling locality had a significant effect when using all four distance metrics,

**TABLE 2** Results of permutational multivariate analysis of variance (PERMANOVA) tests and permutation tests of dispersion on beta distances between gut microbiomes

Distance matrix	Variable	PERMANOVA			Permutation test on dispersion	
		df	R <sup>2</sup>	p	F	p
<i>Full data set (2017–2020)</i>						
Bray–Curtis	Species	14	.047	.162		
	Locality	1	.007	.011*	13.186	.001**
	Year	3	.138	.001**	52.964	.001**
	Sequencing run	1	.130	.001**	171.570	.001**
	Diet type	2	.009	.435		
Jaccard	Species		.053	.001**	4.481	.001**
	Locality		.005	.009**	30.959	.001**
	Year		.080	.001**	151.68	.001**
	Sequencing run		.071	.001**	527.04	.001**
	Diet type		.010	.167		
Unweighted UniFrac	Species	14	.057	.001**	3.189	.001**
	Locality	1	.006	.012*	0.098	.761
	Year	3	.087	.001**	8.033	.001**
	Sequencing run	1	.079	.001**	21.312	.001**
	Diet type	2	.010	.191		
Weighted UniFrac	Species	14	.062	.163		
	Locality	1	.013	.020*	2.155	.146
	Year	3	.016	.146		
	Sequencing run	1	.006	.194		
	Diet type	2	.013	.190		
<i>Batch 1 (2017–2019)</i>						
Bray–Curtis	Species	13	.090	.048*	0.703	.762
	Locality	1	.014	.002**	0.759	.397
	Year	2	.012	.385		
	Diet type	2	.020	.062		
Jaccard	Species		.093	.001**	2.409	.009**
	Locality		.011	.001**	5.607	.020*
	Year		.015	.001**	5.889	.004**
	Diet type		.020	.001**	9.067	.001**
Unweighted UniFrac	Species	13	.103	.001**	3.049	.002**
	Locality	1	.009	.014*	0.002	.971
	Year	2	.015	.034*	2.212	.099
	Diet type	2	.022	.013*	2.161	.118
Weighted UniFrac	Species	13	.085	.380		
	Locality	1	.017	.039*	0.183	.655
	Year	2	.008	.732		
	Diet type	2	.021	.246		

Note: Diet type reflects categorization based on our index of diet specialization (i.e., high diversity, low diversity, intermediate). Asterisks denote significant results: \*\*\* $p < .001$ , \*\* $p < .01$ , \* $p < .05$ .

although the effect size was small (~1%–2% of the variation explained; Table 2). Similarly, year explained a small amount of the variation (~1.5%) when using Jaccard and unweighted UniFrac distances. In the absence of the sequencing batch effect, host species identity accounts for the highest degree of variation in microbiome structure

when using Bray–Curtis (9%,  $p = .048$ ), Jaccard (9.3%,  $p = .001$ ) and unweighted UniFrac distances (10.3%,  $p = .001$ ), generally explaining ~6-fold more of the variation than any other factor considered. Permutation tests indicated that dispersion among species' Jaccard and unweighted UniFrac distances is not homogeneous, which could

account for the significant PERMANOVA result. However, this does not seem to be the case because although dispersion is high for several species causing overlap in PCoA space, species' centroid positions are largely nonoverlapping when using Bray–Curtis, Jaccard and unweighted UniFrac distances (Figure S9), probably reflecting true gut microbiome structuring among species.

In line with our findings of little-to-no correlation between individual diet alpha diversity and gut microbiome alpha diversity, host species diet type did not significantly explain variation between microbiomes in the full data set, nor in the batch 1 subset—with the exception of using Jaccard and unweighted UniFrac distance, in which case diet type explained a small amount of the variation (~2%; Table 2). Dispersion among diet types for Jaccard distance was not homogeneous ( $F = 9.067$ ,  $p = .001$ ), yet diet type centroid positions for Jaccard and unweighted UniFrac distances were nonoverlapping in PCoA space especially for low-diversity diets (Figure S10), indicating some differentiation among gut microbiota for species with more specialized diets.

### 3.4 | Topological congruence analyses

Normalized matching cluster congruence scores for the gut microbiome–host phylogeny topological comparisons were between ~0.4 and 0.8. As congruence scores of zero indicate complete topological congruence, and scores of 1 indicate complete incongruence, these scores reflect intermediate congruences. When analysing all


individuals in the full data set, and within the batch 1 and batch 1-NY subsets, the observed warbler gut microbiome dendrogram was significantly more congruent with the host phylogeny than with randomized dendrograms using Bray–Curtis, Jaccard and weighted UniFrac distances (Table 3, Figure 4a). In the batch 2 subset, the gut microbiome dendrogram was more congruent with the host phylogeny than with randomized dendrograms using Bray–Curtis and unweighted UniFrac distances (Table 3). Thus, the majority of comparisons ( $N = 11$  of 16 comparisons) indicate a positive association between gut microbiome similarity and host phylogenetic relatedness. As Bray–Curtis and weighted UniFrac metrics are weighted by ASV counts, this may indicate that relative abundances of microbial taxa help contribute to the phylogenetic signal in the warbler gut microbiome.

To determine whether gut microbiome similarity better reflects host evolutionary history or host diet similarity, we repeated the topological analyses above instead using a dendrogram clustered from the diet OTU distances in place of the host phylogeny (Table 3). Among comparisons, congruence scores were generally lower (indicating better congruence) for microbiome–host phylogeny comparisons than for microbiome–diet comparisons (Figure S11), although there are some exceptions. Importantly, only six of 16 microbiome–diet comparisons were significantly more congruent than random. Three of these comparisons were of Jaccard distance, which only considers ASV presence/absence. Further, in two other instances both considering Bray–Curtis distances, congruence scores for the microbiome–host phylogeny comparison were lower than for

**TABLE 3** Summary of topological congruences between species-level gut microbiome dendrograms and the host phylogeny (left), and between species-level diet dendrograms (right), and of individual-level Mantel tests

	N spp.	Distance metric	Microbiome–host phylogeny		Microbiome–diet	
			Matching cluster congruence score	Mantel $r$	Matching cluster congruence scores	Mantel $r$
Full data set (2017–2020)	15	Bray–Curtis	0.52***	.02	0.57	.06*
		Jaccard	0.58**	.04	0.56***	.17**
		Unweighted UniFrac	0.68	.10**	0.74	.16**
		Weighted UniFrac	0.58**	.01	0.51**	.03
Batch 1 (2017–2019)	14	Bray–Curtis	0.45***	.09*	0.70	.09*
		Jaccard	0.45***	.18**	0.39***	.21**
		Unweighted UniFrac	0.71	.19**	0.64	.22**
		Weighted UniFrac	0.57**	–.003	0.61	–.05
Batch 1-NY (2017–2019)	13	Bray–Curtis	0.44***	.15**	0.53*	.11*
		Jaccard	0.52**	.23**	0.44***	.23**
		Unweighted UniFrac	0.73	.27**	0.69	.25**
		Weighted UniFrac	0.52**	.07	0.74	–.09
Batch 2 (2020)	14	Bray–Curtis	0.56**	.07	0.62*	.01
		Jaccard	0.72	.03	0.67	–.03
		Unweighted UniFrac	0.59*	.04	0.64	–.02
		Weighted UniFrac	0.79	–.02	0.73	.06

Note: N spp. = number of species analysed, and matching cluster congruence scores are normalized where 0 = complete congruence and 1 = complete incongruence. Asterisks denote significant results: \*\*\* $p < .001$ , \*\* $p < .01$ , \* $p < .05$ .



**FIGURE 4** Summary of phylosymbiosis analyses. (a) Topological congruence analyses of the association between the gut microbiome and host phylogeny (left), the gut microbiome and diet (middle), and the diet and host phylogeny (right). Microbiome and diet dendrograms were constructed using Bray–Curtis distances of mean within-species ASV/OTU counts. Matching cluster congruence scores are normalized where 0 = complete congruence, and 1 = complete incongruence. See Table 1 for species codes. (b) Scatter plots of individual-level microbiome vs. host evolutionary distances (left), microbiome vs. diet distances (middle), and diet vs. host evolutionary distances. Diet and microbiome distances are of the Bray–Curtis metric. [Colour figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1111/mec.16762)]

the microbiome–diet comparison (batch 2, batch 1-NY; Table 3). Collectively, these results suggest a closer association between gut microbiome structure and host evolutionary history than with host diet.

Finally, we examined the association between the host phylogeny and diet dendrograms from the batch 1 subset to determine whether the significant associations we detected between the gut microbiome and diet could be due to a phylogenetic signal of the diet. For all four distance metrics, the diet–host phylogeny comparison was significantly more congruent than random (Bray–Curtis normalized matching cluster score = 0.56,  $p = .007$ , Jaccard normalized matching cluster score = 0.52,  $p = .003$ , unweighted UniFrac normalized matching cluster score = 0.51,  $p = .001$ , weighted UniFrac normalized matching cluster score = 0.57,  $p = .017$ ). These scores reflect intermediate congruence between the diet dendrogram and host phylogeny, and are similar but slightly higher (less congruent) on average than significant congruence scores between the microbiome dendrogram and host phylogeny in batch 1 (Figure 4a, Table 3).

### 3.5 | Mantel tests

Mantel tests indicated a positive relationship between individual-level microbiome distances and pairwise evolutionary distances (Mantel  $r = .09$ – $.27$ ) in the batch 1 and batch 1-NY data sets for Bray–Curtis, Jaccard and unweighted UniFrac distances, and in the full data set for unweighted UniFrac distance (Table 3, Figure 4b). Mantel tests also indicated a positive relationship between individual-level microbiome distances and diet distances (Mantel  $r = .06$ – $.25$ ) in the

full data set, batch 1 and batch 1-NY subsets using Bray–Curtis, Jaccard and unweighted UniFrac distances (Table 3, Figure 4b).

We also tested the relationship between diet matrices and pairwise evolutionary distances in the batch 1 subset, and found a positive association for Jaccard diet distance (Mantel  $r = .10$ ,  $p = .027$ ), and unweighted UniFrac diet distance (Mantel  $r = .13$ ,  $p = .004$ ). Notably, across all Mantel tests, most significant correlations were detected when using unweighted distance matrices (Jaccard and unweighted UniFrac).

## 4 | DISCUSSION

We performed faecal metabarcoding to examine environmental and evolutionary influences on gut microbiome structure in breeding wood-warblers. Our analyses collectively support host taxonomy as the strongest driver of gut microbiome structure while environmental factors, including diet type, showed lesser effects. At the individual level, diet alpha diversity showed little to no association with microbiome alpha diversity. Further, on average, more closely related species tended to harbour more similar gut microbiomes, and gut microbiome similarity was less closely associated with diet similarity, suggesting host evolutionary history may play a large role in shaping host–microbe interactions in this clade. We also detected a relatively strong batch effect of sequencing run on gut microbiome diversity, and by analysing within-batch subsets of our data we saw that this had obscured the signal of the biological factors we considered in our analyses. Thus, these results highlight caution for other researchers about whether or not to divide samples across

sequencing lanes and this should be a serious consideration in future metabarcoding studies.

#### 4.1 | The wood-warbler gut microbiome

Wood-warbler gut microbiomes were dominated by Proteobacteria and Firmicutes, which is consistent with other studies of other free-living passerines (e.g., Bodawatta, Koane, et al., 2021; Hird et al., 2015). The most prevalent ASV, a Proteobacteria in the family Yersiniaceae, was observed in ~60% of individuals and occurred in all host species examined, but only a very small proportion of ASVs were represented in >30% of the individuals sequenced. These results may reflect a shared signature of the passerine gut microbiome in wood-warblers at higher taxonomic levels, yet a high level of variability among individuals, especially for lower abundance taxa.

The most dominant bacterial phyla in the current study were also identified as highly abundant in the only migratory cycle study of recaptured warblers to date, which focused on Kirtland's warblers (*Setophaga kirtlandii*; Skeen et al., 2021), a species that does not breed in our study areas. Although arrival on the breeding grounds was accompanied by a shift from a Kirtland's warbler gut microbiome dominated by Firmicutes to one dominated by Proteobacteria, both phyla were highly abundant across the migratory cycle. The most prevalent taxonomic classes in the current study (Gammaproteobacteria, Alphaproteobacteria and Bacilli) also dominated gut microbiomes of breeding Kirtland's warblers (Skeen et al., 2021). However, Clostridia was one of the most abundant taxa in Kirtland's warblers but was found at low prevalence among individuals in the current study and made up only <2% of the total reads sequenced. This may suggest that Kirtland's warblers, a near-threatened Caribbean migrant with highly specialized habitat requirements, differ in gut microbiome structure from closely related parulids breeding nearby. This differentiation would be consistent with our findings of a relatively strong role of host taxonomy and evolutionary history, and/or associated environmental factors that we were unable to resolve with our data set, in shaping the parulid gut microbiome (see below).

In this study, sampling locality consistently explained 1%–2% of the variation between microbiomes across data sets and distance metrics considered. Samples were collected from two forested localities in Eastern North America roughly 400km apart, a distance that is probably not large enough to generate significant population genetic structure within warbler host species due to a lack of potential barriers to gene flow (e.g., yellow-rumped warblers, *Setophaga coronata*; Toews et al., 2016). However, our results suggest this distance may be sufficient in scale to affect subtle changes in gut microbe communities. Interestingly, the amount of variation explained by sampling locality here is similar to that reported in other passerine studies (San Juan et al., 2021; Teyssier et al., 2020), despite this study encompassing a larger geographical area. For example, habitat type explained ~4% of the variation between passerine microbiomes within a 43-km agricultural study area in Costa Rica (San Juan et al., 2021), suggesting habitat features may be more important

than geographical distance between sites. Although we did not include habitat features as a factor in our analyses, notable differences between our study sites include an abundance of *Rhododendron* (*R. maximum*) and mountain laurel (*Kalmia latifolia*) in the understory at our Pennsylvania localities, whereas these shrubs do not occur in our New York localities. This and other habitat differences could conceivably contribute to the differences we observed in gut microbiota between our sites.

When analysing a subset of samples from a single sequencing run, sampling year explained a similar proportion of variation between microbiomes as did sampling locality, but tended not to be significant. This may indicate that wood-warbler microbiomes are stable across breeding seasons, despite annual long-distance longitudinal migration to and from tropical nonbreeding grounds, which is probably associated with changes in foraging strategies. This is consistent with other passerine studies that found no difference in gut microbiome diversity across consecutive breeding seasons (Benskin et al., 2015; Escallón et al., 2019), but it is important to note that in our data set, each year represents a different cohort of individuals. In migratory species, it will be desirable to resample the same individuals on the nonbreeding and breeding grounds across multiple cycles to disentangle temporal effects from those of habitat, diet and geographical locality (Bodawatta, Hird, et al., 2021; Skeen et al., 2021).

#### 4.2 | Diet diversity is not tightly linked to gut microbiome diversity in wood-warblers

By sequencing both arthropod COI gene and bacterial 16S rRNA gene metabarcoding libraries from the same faecal samples, we were able to directly examine the relationship between natural diet diversity and gut microbiome diversity of the same individuals. Our strategy revealed that when analysing three different metrics of within-individual (alpha) diversity, diet diversity was not correlated with microbiome diversity with the exception of a weak correlation in the batch 1 data when using the Chao1 index, which is neither phylogenetically aware nor weighted by ASV/OTU abundance (Figure 2a). Although individuals of species with low-diversity diets tended to have reduced gut microbiome alpha diversity and individuals of species with high-diversity diets tended to have increased microbiome alpha diversity, this pattern was not significant (Figure 2c). Further, when looking at between-individual (beta) diversity, diet type explained only ~2% of the variation between individuals and only when using unweighted distance metrics. In this case, individuals of species with more specialized (less diverse) diets tended to drive this pattern (Figure S10). This provides some evidence that diet richness may be weakly associated with gut microbiome richness, although we were unable to detect significant associations with these analyses when using our full data set, which may be due to the batch effect. Similarly, Mantel tests provided some support for a positive relationship between individual gut microbiome similarity and diet similarity especially for unweighted distance metrics, but

the association is not strong (Table 3). Thus, in contrast to our prediction, diversity of the diet generally did not explain variation in the gut microbiome. This may suggest a high-diversity diet either does not generally provide wood-warblers with an increased availability of potential gut colonists, or gut microbe colonization is not strongly affected by diet diversity. Similarly, in a study of two species of freshwater fish, Bolnick et al. (2014) found the relationship between diet diversity and gut microbiome diversity was not linear and fish with a specialized diet actually harboured a more diverse gut microbiome.

Despite our finding of a weak relationship between diet diversity and gut microbiome diversity, many studies have shown host diet indeed influences the avian gut microbiome. Broad categorization of natural feeding guild and diet type explain differences in the gut microbiomes of wild passerines in New Guinea and of zoo and farm birds in China, respectively (Bodawatta, Koane, et al., 2021; Xiao et al., 2021). Further, experimental manipulations of passerine diets have been associated with shifts in gut microbiome diversity and composition (Davidson et al., 2020; Knutie, 2020; Pekarsky et al., 2021; Teyssier et al., 2020). In the current study, we analysed natural diets of breeding wood-warblers, which are known to primarily eat insects (Birds of the World, 2022; MacArthur, 1958). Our metabarcoding results indicate a substantial portion of the diet is also arachnid-based. However, diet alpha diversity did not differ among species, and relative proportions of arthropod classes in the diet were similar (Figure 1c). The lack of species with a highly specialized diet (at the scale analysed here) that were included in this study may make wood-warblers a poor system for untangling the effect of diet diversity on gut microbiome diversity, and future dual diet-microbiome metabarcoding studies could also include birds with clear distinctions in dietary guild for comparison (e.g., extreme diet specialists, aerial insectivores). We note that we did not consider fine-scale spatial partitioning of the feeding niche as an explanatory variable in this study, something wood-warblers are well known for (MacArthur, 1958). Further, it is possible that because we examined broad-scale patterns in diet diversity at the OTU level, we were not able to identify components of the diet (e.g., nutritional values of arthropods) that possibly underlie gut microbiome structure. We also note that although wood-warblers are primarily insectivores, some species are known to supplement their diet with fruit, especially in the nonbreeding season (Birds of the World, 2022). Our study design did not allow us to examine effects of any nonarthropod components of the diet, which may influence gut microbiota (Bodawatta et al., 2022). Nevertheless, our results suggest dietary arthropod diversity does not scale directly with gut microbiome diversity in breeding wood-warblers.

### 4.3 | Host evolution as the main driver of wood-warbler gut microbiome structure

Among the biological factors considered in this study, host species stands out as the variable that explains the largest amount of

variation between microbiomes. Further, species-level 16S dendrograms were generally more concordant with the host phylogeny than with COI dendrograms (Figure 4a, Table 3). We also found the host phylogeny to be concordant with COI diet dendrograms, suggesting the weaker associations we did detect between the diet and gut microbiome may have arisen due to a phylogenetic signal of both the diet (Miller et al., 2022) and microbiome. Together with our findings of little environmental influence on the wood-warbler microbiome, this may suggest that host evolutionary history rather than differences in species' ecological niche is the main driver of microbiome differentiation between wood-warbler species.

Mantel analyses of individual-level matrices revealed a somewhat contrasting pattern, showing support for positive associations between the gut microbiome and evolutionary distance and a similar level of support for a positive relationship between the gut microbiome and diet distance. Similar to the topological congruence analysis, these analyses also showed some support for a relationship between the diet and evolutionary distance. In these analyses, most of the significant associations involving the diet arose using unweighted distance metrics. These results are consistent with our other diet diversity analyses by suggesting community richness is driving these patterns.

The conflicting pattern revealed by the topological congruence analyses and Mantel tests may be explained by at least two factors. First, although they are complementary tests of phyllosymbiosis, topological congruence analyses and Mantel tests fundamentally rely on different information. Topological congruence analyses do not rely on branch lengths or directly consider evolutionary or beta distances, whereas Mantel tests measure the correlation between two distance matrices. Because changes in microbiome community structure may be much more rapid than evolutionary changes between host genomes, topological congruence analyses may be a more conservative test of phyllosymbiosis (Lim & Bordenstein, 2020).

Second, we used species-averaged ASV/OTU counts in the topological congruence analyses in order to summarize variation within each species, whereas our Mantel tests were of distance matrices based on individual-level data. Across all of our analyses of individual-level data, both alpha and beta diversity of the diet and microbiome were quite variable, even within species. For example, Bray-Curtis distances between individuals of the same species ranged from ~0.07 to 1 for the gut microbiome, and from ~0.25 to 1 for the diet (Figure 4b). This may suggest that the high level of variation within species obscured phylogenetic signal in gut microbiome and diet similarities at the individual level in Mantel tests.

Host species identity was the biological factor that explained the highest degree of variation between microbiota (Table 2), suggesting the mean ASV counts used in topological congruence analyses may capture unique features within host species. Collectively, our analyses support a tighter association between the gut microbiome and host evolutionary history than between the gut microbiome and diet similarity when looking at the level of host species. This phylogenetic signal of gut microbiome structure is well supported in nonflying mammals and insects, but has been less well supported in birds. Avian

gut microbiome studies generally support differences between host species (Hird et al., 2015; San Juan et al., 2020; Capunitan et al., 2020; but see Hird et al., 2014), but phylosymbiosis was not supported in New Guinean passerines (Bodawatta, Koane, et al., 2021) and the signal was weak among captive cranes and in two passerine studies (Kropáčková et al., 2017; Loo et al., 2019; Trevelline et al., 2020). In the current study, concordance between the wood-warbler phylogeny and gut microbiome dendrogram was moderate and similar to that reported for cranes in captivity (Trevelline et al., 2020) and passerines in the Czech Republic (Kropáčková et al., 2017). Thus, our results support the view that phylosymbiosis is weaker in birds than in mammals (Song et al., 2020; Youngblut et al., 2019) and uniquely demonstrate that in wood-warblers, a stronger influence of diet (as measured by species-level diet diversity) does not account for this discrepancy. Our findings of high variability of gut microbiomes for individuals within the same species may explain the lack of a consensus about phylosymbiosis in the avian literature, and particularly among studies that analysed fewer individuals per species.

A phylogenetically conserved gut microbiome may provide the opportunity for co-adaptation between hosts and their gut microbes, which could implicate microbiomes in complex host evolutionary processes, including speciation (Brucker & Bordenstein, 2012). Long-term co-evolution between hosts and microbiota could explain phylosymbiosis, but this pattern could also arise under ecological filtering. Mazel et al. (2018) used simulations to show that under ecological filtering, the strength of phylosymbiosis is determined by the strength of the phylogenetic signal in the host trait underlying microbe colonization. It has been hypothesized that convergence of bat and avian gut microbiomes is due to reduced gut length, an adaptation to powered flight, which may favour rapid turnover in gut microbiota, thus accounting for the weakened phylogenetic signal in avian gut microbiomes compared to nonflying mammals (Song et al., 2020). Consistent with this, Bodawatta, Koane, et al. (2021) found a negative association between passerine body mass—a proxy for gut length—and both gut microbiome richness and divergence. This might lead to the prediction that phylogenetic signal of the gut microbiome should be strongest in large-bodied birds and weakest in small-bodied birds. However, the current data do not support this, as is highlighted by the results presented here. The strength of phylosymbiosis reported here for wood-warblers—small-bodied species weighing ~6–20g—is similar to that reported for cranes (Trevelline et al., 2020), which are several hundred times heavier. Thus, additional study is necessary to elucidate the effect of gut retention time on gut microbiome structure, and of other phylogenetically conserved avian traits or habitat preferences, including diet, that may mediate the colonization and maintenance of gut microbiomes.

Further study is also necessary to understand the biological relevance of taxonomic differences and of phylogenetic signal in gut microbiome structure between hosts. Experimental studies have shown antibiotic treatment administered to nestlings results in faster growth rates (Coates et al., 1963; Kohl et al., 2018; Potti et al., 2002), and caeca of germ-free chickens exhibit altered gene expression and notably do not express immunoglobulins (Volf et al., 2017). Thus, it

is clear that gut microbiota impose constraints on host development and immune function, but how species-differences in natural gut microbiota composition might impact host fitness is unknown. It is important to note that although we observed an effect of host taxonomy on gut microbiome structure, this does not necessarily imply functional differences in gut microbiota between hosts. However, due to microbiome differentiation between host species we may predict disruption of these communities for admixed individuals upon hybridization (Brucker & Bordenstein, 2012). Wood-warblers are well known to hybridize and occasionally even form intergeneric hybrids (Toews et al., 2018, 2020), making this clade an excellent system that can be used to tease apart evolutionary from ecological influences on the gut microbiome as well as the potential role of the microbiome in hybrid dysfunction.

#### 4.4 | Limitations of metabarcoding and caution against sequencing batch effects

Our study faces several limitations inherent to metabarcoding approaches (Taberlet et al., 2012). First, we used primers to target gut bacteria and arthropod taxa. However, gut communities are very complex and also include archaea, fungi, nonarthropod diet taxa (discussed above) and other eukaryotes including gut parasites. These other taxa probably influence gut microbiome dynamics, which we were not able to consider here. Further, we were limited by taxonomic databases to identify gut bacteria and dietary arthropods. A small proportion of the taxa analysed here could not be identified beyond domain, including two ASVs in the core warbler gut microbiome (Tables S3 and S4). Finally, PCR steps introduce bias in the detection of taxa and estimates of their abundance (James et al., 2022). We took steps to minimize the effects of bias by sequencing multiple individuals per host species, and we analysed mean within-species OTU counts in certain analyses. This approach should help limit the impact of artificially biased abundances when analysing differences among host species, as artificial bias is not expected to be consistent across individuals. We also randomized the order in which we processed samples during library preparation, such that each plate contained samples collected across multiple species, years and localities in case of plate-specific PCR batch effects. To further limit PCR bias in 16S libraries, we included three technical replicates for each individual and pooled replicates before sequencing (Caporaso et al., 2011). However, for COI libraries, we only sequenced a single PCR per individual. This approach probably led us to underestimate the richness of diet taxa, and we probably were not able to detect rare diet taxa (Alberdi et al., 2018).

A strength of our study is that we were able to include large sample sizes, although we split our sequencing into two batches. We prepared and sequenced our 16S and COI libraries in two different batches and the resulting yields were quite different (Figure S5). This strategy was desirable because it allowed us to process samples as they became available and it increased our overall sample sizes. However, we found the technical artefacts this introduced

were not trivial (Figure S8), and similar to batch effects in other studies (Gibbons et al., 2018; Lou & Therkildsen, 2022) it obscured the signal of the biological factors we tested (Table 2). Our topological congruence analysis seemed to be robust to the batch effect as our results were similar across data sets, although it is possible that batch effects obscured the signal of phyllosymbiosis in previous avian gut microbiome studies. It is possible that batch effects are less of a concern for mammalian and other systems, where the signal of host phylogeny on gut microbiome structure is stronger than in birds. Similarly, the batch effect was less strong in the COI data, which we also processed and sequenced in two batches (37% of total arthropod OTUs detected in both batches compared to 6% for bacterial ASVs). This may be due to a more rapid saturation of the accumulation curve for arthropod taxa than for bacterial taxa that are present in the warbler gut (Figure S2).

Future methodological study of the consequences of batch effects in metabarcoding studies is warranted. We recommend that metabarcoding studies report sequences of technical replicates (PCRs amplified from the same sample within a batch) and positives (same sample sequenced across batches) which may help clarify when it is appropriate to make direct comparison of data sequenced in different batches.

## 5 | CONCLUDING REMARKS

Our data highlight many outstanding questions regarding avian microbiomes and the ongoing need to characterize microbiomes of wild birds (Hird, 2017). Wood-warbler gut microbiomes are dominated by Proteobacteria and Firmicutes, and on average, closely related host species share more similar gut microbiomes. We found little influence of sampling year, geographical locality or diet diversity on gut microbiome structure, and thus the majority of the variation between microbiota was left unexplained. Our results may suggest the phylogenetic signal in gut microbiome structure is tied more closely to host traits than to host environment, yet the mechanisms driving this signal and possible functional consequences for hosts are not clear.

The level of phylogenetic signal in gut microbiome structure we detected is similar to that detected for larger-bodied birds (Trevelline et al., 2020), suggesting small body size does not preclude phyllosymbiosis. Further study is necessary to understand the relationship between host body size, gut retention time and gut microbe colonization. Although we found that broad-scale measures of diet diversity are not closely related to gut microbiome diversity, future studies should explore how components of the diet (e.g., dominant arthropod taxa, energetic values of food items) might influence the gut microbiome, including by way of their influence on host traits (e.g., gut pH). Wood-warblers represent a promising system to continue addressing outstanding ecological and evolutionary questions about the avian microbiome, including how microbiomes may influence and respond to adaptive radiation (Bodawatta, Hird, et al., 2021).

## AUTHOR CONTRIBUTIONS

MDB and DPLT designed the study with input from all authors. DPLT, ETM, MDB and AWW collected samples, and AWW processed samples and prepared sequencing libraries. All authors took part in data analysis. MDB and ABC wrote the manuscript, and all authors revised and approved the final version.

## ACKNOWLEDGEMENTS

We thank Brian Trevelline and Emily Davenport for helpful discussion and advice on data analyses, and three anonymous reviewers for feedback that greatly improved our manuscript. We also thank Laura Porturas and Lan-Nhi Phung for assistance in the field. This material is based upon work supported by the NSF Postdoctoral Research Fellowships in Biology Program under Grant No. 2010679. This work was also supported by the Department of Biology and the Huck Institute for Life Sciences at Penn State University, and by the Cornell Lab of Ornithology.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## DATA AVAILABILITY STATEMENT

The sequencing data analyzed in this study is available at the NCBI Short Read Archive under BioProject PRJNA889412. Associated scripts and metadata are available on GitHub at [https://github.com/baizm/parulid\\_microbiome](https://github.com/baizm/parulid_microbiome) and on Dryad at <https://doi.org/10.5061/dryad.905qfttps>.

## BENEFITS-SHARING

Benefits from this research accrue from the sharing of our data on public resources as described above.

## ORCID

Marcella D. Baiz  <https://orcid.org/0000-0002-1629-6737>

David P. L. Toews  <https://orcid.org/0000-0002-9763-0476>

## REFERENCES

- Alberdi, A., Aizpurua, O., Gilbert, M. T. P., & Bohmann, K. (2018). Scrutinizing key steps for reliable metabarcoding of environmental samples. *Methods in Ecology and Evolution*, 9(1), 134–147. <https://doi.org/10.1111/2041-210X.12849>
- Baiz, M. D., Wood, A. W., Brelford, A., Lovette, I. J., & Toews, D. P. (2021). Pigmentation genes show evidence of repeated divergence and multiple bouts of introgression in Setophaga warblers. *Current Biology*, 31(3), 643–649. <https://doi.org/10.1016/j.cub.2020.10.094>
- Barker, F. K., Burns, K. J., Klicka, J., Lanyon, S. M., & Lovette, I. J. (2015). New insights into New World biogeography: An integrated view from the phylogeny of blackbirds, cardinals, sparrows, tanagers, warblers, and allies. *The Auk: Ornithological Advances*, 132(2), 333–348. <https://doi.org/10.1642/AUK-14-110.1>
- Battaglia, T. (2022). *Btools: A suite of R function for all types of microbial diversity analyses*. R package version 0.0.1.
- Benskin, C. M. H., Rhodes, G., Pickup, R. W., Mainwaring, M. C., Wilson, K., & Hartley, I. R. (2015). Life history correlates of fecal bacterial species richness in a wild population of the blue tit *Cyanistes*



- caeruleus. *Ecology and Evolution*, 5(4), 821–835. <https://doi.org/10.1002/ece3.1384>
- Birds of the World (2022). In S. M. Billerman, B. K. Keeney, P. G. Rodewald, & T. S. Schulenberg (Eds.), *Birds of the World*. Cornell Laboratory of Ornithology. <https://birdsoftheworld.org/bow/home>
- Bodawatta, K. H., Hird, S. M., Grond, K., Poulsen, M., & Jønsson, K. A. (2021). Avian gut microbiomes taking flight. *Trends in Microbiology*, 30, 268–280. <https://doi.org/10.1016/j.tim.2021.07.003>
- Bodawatta, K. H., Klečková, I., Klečka, J., Pužejová, K., Koane, B., Poulsen, M., Jønsson, K. A., & Sam, K. (2022). Specific gut bacterial responses to natural diets of tropical birds. *Scientific Reports*, 12(1), 1–15. <https://doi.org/10.1038/s41598-022-04808-9>
- Bodawatta, K. H., Koane, B., Maiah, G., Sam, K., Poulsen, M., & Jønsson, K. A. (2021). Species-specific but not phyllosymbiotic gut microbiomes of new Guinean passerine birds are shaped by diet and flight-associated gut modifications. *Proceedings of the Royal Society B*, 288(1949), 20210446. <https://doi.org/10.1098/rspb.2021.0446>
- Bogdanowicz, D., & Giaro, K. (2013). On a matching distance between rooted phylogenetic trees. *International Journal of Applied Mathematics and Computer Science*, 23(3), 669–684. <https://doi.org/10.2478/amcs-2013-0050>
- Bogdanowicz, D., Giaro, K., & Wróbel, B. (2012). TreeCmp: Comparison of trees in polynomial time. *Evolutionary Bioinformatics*, 8, EBO-S9657. <https://doi.org/10.4137/EBO.S9657>
- Bolnick, D. I., Snowberg, L. K., Hirsch, P. E., Lauber, C. L., Knight, R., Caporaso, J. G., & Svanbäck, R. (2014). Individuals' diet diversity influences gut microbial diversity in two freshwater fish (threespine stickleback and Eurasian perch). *Ecology Letters*, 17(8), 979–987. <https://doi.org/10.1111/ele.12301>
- Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., al-Ghalith, G. A., Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K., Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., ... Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Bio/technology (Nature Publishing Company)*, 37(8), 852–857. <https://doi.org/10.1038/s41587-019-0209-9>
- Brooks, A. W., Kohl, K. D., Brucker, R. M., van Opstal, E. J., & Bordenstein, S. R. (2016). Phyllosymbiosis: Relationships and functional effects of microbial communities across host evolutionary history. *PLoS Biology*, 14(11), e2000225. <https://doi.org/10.1371/journal.pbio.1002587>
- Brucker, R. M., & Bordenstein, S. R. (2012). Speciation by symbiosis. *Trends in Ecology & Evolution*, 27(8), 443–451. <https://doi.org/10.1016/j.tree.2012.03.011>
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S. M., Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J. A., Smith, G., & Knight, R. (2012). Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *The ISME Journal*, 6(8), 1621–1624. <https://doi.org/10.1038/ismej.2012.8>
- Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Lozupone, C. A., Turnbaugh, P. J., Fierer, N., & Knight, R. (2011). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample. *Proceedings of the National Academy of Sciences*, 108(supplement\_1), 4516–4522. <https://doi.org/10.1073/pnas.1000080107>
- Capunitan, D. C., Johnson, O., Terrill, R. S., & Hird, S. M. (2020). Evolutionary signal in the gut microbiomes of 74 bird species from Equatorial Guinea. *Molecular Ecology*, 29(4), 829–847. <https://doi.org/10.1111/mec.15354>
- Coates, M. E., Fuller, R., Harrison, G. F., Lev, M., & Suffolk, S. F. (1963). A comparison of the growth of chicks in the Gustafsson germ-free apparatus and in a conventional environment, with and without dietary supplements of penicillin. *British Journal of Nutrition*, 17(1), 141–150. <https://doi.org/10.1079/BJN19630015>
- Davidson, G. L., Wiley, N., Cooke, A. C., Johnson, C. N., Fouhy, F., Reichert, M. S., de la Hera, I., Crane, J. M. S., Kulahci, I. G., Ross, R. P., Stanton, C., & Quinn, J. L. (2020). Diet induces parallel changes to the gut microbiota and problem solving performance in a wild bird. *Scientific reports.*, 10(1), 20783. <https://doi.org/10.1038/s41598-020-77256-y>
- Davis, N. M., Proctor, D. M., Holmes, S. P., Relman, D. A., & Callahan, B. J. (2018). Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. *Microbiome*, 6(1), 226. <https://doi.org/10.1186/s40168-018-0605-2>
- Escallón, C., Belden, L. K., & Moore, I. T. (2019). The cloacal microbiome changes with the breeding season in a wild bird. *Integrative Organismal Biology*, 1(1), oby009. <https://doi.org/10.1093/iob/oby009>
- Gibbons, S. M., Duvallet, C., & Alm, E. J. (2018). Correcting for batch effects in case-control microbiome studies. *PLoS Computational Biology*, 14(4), e1006102. <https://doi.org/10.1371/journal.pcbi.1006102>
- Grond, K., Santo Domingo, J. W., Lanctot, R. B., Jumpponen, A., Bentzen, R. L., Boldenow, M. L., Brown, S. C., Casler, B., Cunningham, J. A., Doll, A. C., Freeman, S., Hill, B. L., Kendall, S. J., Kwon, E., Liebezeit, J. R., Pirie-Dominix, L., Rausch, J., & Sandercock, B. K. (2019). Composition and drivers of gut microbial communities in Arctic-breeding shorebirds. *Frontiers in virology*, 10, 2258. <https://doi.org/10.3389/fmicb.2019.02258>
- Hird, S. M. (2017). Evolutionary biology needs wild microbiomes. *Frontiers in Microbiology*, 8, 725. <https://doi.org/10.3389/fmicb.2017.00725>
- Hird, S. M., Carstens, B. C., Cardiff, S. W., Dittmann, D. L., & Brumfield, R. T. (2014). Sampling locality is more detectable than taxonomy or ecology in the gut microbiota of the brood-parasitic Brown-headed cowbird (*Molothrus ater*). *PeerJ*, 2, e321. <https://doi.org/10.7717/peerj.321>
- Hird, S. M., Sánchez, C., Carstens, B. C., & Brumfield, R. T. (2015). Comparative gut microbiota of 59 neotropical bird species. *Frontiers in Microbiology*, 6, 1403. <https://doi.org/10.3389/fmicb.2015.01403>
- Hughes, J. B., & Hellmann, J. J. (2005). The application of rarefaction techniques to molecular inventories of microbial diversity. *Methods in Enzymology*, 397, 292–308. [https://doi.org/10.1016/S0076-6879\(05\)97017-1](https://doi.org/10.1016/S0076-6879(05)97017-1)
- James, A. R., Geber, M. A., & Toews, D. P. (2022). Molecular assays of pollen use consistently reflect pollinator visitation patterns in a system of flowering plants. *Molecular Ecology Resources*, 22(1), 361–374. <https://doi.org/10.1111/1755-0998.13468>
- Jusino, M. A., Banik, M. T., Palmer, J. M., Wray, A. K., Xiao, L., Pelton, E., Barber, J. R., Kawahara, A. Y., Gratton, C., Peery, M. Z., & Lindner, D. L. (2019). An improved method for utilizing high-throughput amplicon sequencing to determine the diets of insectivorous animals. *Molecular Ecology Resources*, 19(1), 176–190. <https://doi.org/10.1111/1755-0998.12951>
- Knutie, S. A. (2020). Food supplementation affects gut microbiota and immunological resistance to parasites in a wild bird species. *Journal of Applied Ecology*, 57(3), 536–547. <https://doi.org/10.1111/1365-2664.13567>
- Kohl, K. D., Brun, A., Bordenstein, S. R., Caviedes-Vidal, E., & Karasov, W. H. (2018). Gut microbes limit growth in house sparrow nestlings (*Passer domesticus*) but not through limitations in digestive capacity. *Integrative Zoology*, 13(2), 139–151. <https://doi.org/10.1111/1749-4877.12289>
- Kropáčková, L., Těšický, M., Albrecht, T., Kubovčík, J., Čížková, D., Tomášek, O., Martin, J., Bobek, L., Králová, T., Procházka, P., & Kreisinger, J. (2017). Codiversification of gastrointestinal microbiota and phylogeny in passerines is not explained by ecological divergence. *Molecular Ecology*, 26(19), 5292–5304. <https://doi.org/10.1111/mec.14144>

- Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: A resource for timelines, timetrees, and divergence times. *Molecular Biology and Evolution*, 34(7), 1812–1819. <https://doi.org/10.1093/molbev/msx116>
- Lim, S. J., & Bordenstein, S. R. (2020). An introduction to phyllosymbiosis. *Proceedings of the Royal Society B*, 287(1922), 20192900. <https://doi.org/10.1098/rspb.2019.2900>
- Loo, W. T., García-Loor, J., Dudaniec, R. Y., Kleindorfer, S., & Cavanaugh, C. M. (2019). Host phylogeny, diet, and habitat differentiate the gut microbiomes of Darwin's finches on Santa Cruz Island. *Scientific Reports*, 9(1), 18781. <https://doi.org/10.1038/s41598-019-54869-6>
- Lou, R. N., & Therikildsen, N. O. (2022). Batch effects in population genomic studies with low-coverage whole genome sequencing data: Causes, detection and mitigation. *Molecular Ecology Resources*, 22(5), 1678–1692. <https://doi.org/10.1111/1755-0998.13559>
- Lovette, I. J., Pérez-Emán, J. L., Sullivan, J. P., Banks, R. C., Fiorentino, I., Córdoba-Córdoba, S., Echeverry-Galvis, M., Barker, F. K., Burns, K. J., Klicka, J., Lanyon, S. M., & Bermingham, E. (2010). A comprehensive multilocus phylogeny for the wood-warblers and a revised classification of the Parulidae (Aves). *Molecular Phylogenetics and Evolution*, 57(2), 753–770. <https://doi.org/10.1016/j.ympev.2010.07.018>
- MacArthur, R. H. (1958). Population ecology of some warblers of north-eastern coniferous forests. *Ecology*, 39(4), 599–619. <https://doi.org/10.2307/1931600>
- Mazel, F., Davis, K. M., Loudon, A., Kwong, W. K., Groussin, M., & Parfrey, L. W. (2018). Is host filtering the main driver of phyllosymbiosis across the tree of life? *Msystems*, 3(5), e00097-18. <https://doi.org/10.1128/mSystems.00097-18>
- McMurdie, P. J., & Holmes, S. (2013). Phyloseq: An R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS One*, 8(4), e61217. <https://doi.org/10.1371/journal.pone.0061217>
- Michel, A. J., Ward, L. M., Goffredi, S. K., Dawson, K. S., Baldassarre, D. T., Brenner, A., Gotanda, K. M., McCormack, J. E., Mullin, S. W., O'Neill, A., Tender, G. S., Uy, J. A. C., Yu, K., Orphan, V. J., & Chaves, J. A. (2018). The gut of the finch: Uniqueness of the gut microbiome of the Galápagos vampire finch. *Microbiome*, 6(1), 1–14. <https://doi.org/10.1186/s40168-018-0555-8>
- Miller, E. T., Wood, A., Baiz, M. D., Welch, A. J., Fleischer, R. C., Dale, A. S., & Toews, D. P. (2022). Re-assessing niche partitioning in MacArthur's warblers: Foraging behavior, morphology, and diet metabarcoding in a phylogenetic context. *bioRxiv*. <https://doi.org/10.1101/2022.08.26.505503>
- Oksanen, J., Guillaume Blanchet, F., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Peter, S., Henry, M., Stevens, H., Szocs, E., & Wagner, H. (2020). *Vegan: Community Ecology Package. R package version 2.5-7*.
- Pekarsky, S., Corl, A., Turjeman, S., Kamath, P. L., Getz, W. M., Bowie, R. C., Markin, Y., & Nathan, R. (2021). Drivers of change and stability in the gut microbiota of an omnivorous avian migrant exposed to artificial food supplementation. *Molecular Ecology*, 30(19), 4723–4739. <https://doi.org/10.1111/mec.16079>
- Potti, J., Moreno, J., Yorio, P., Briones, V., García-Borboroglu, P., Villar, S., & Ballesteros, C. (2002). Bacteria divert resources from growth for magellanic penguin chicks. *Ecology Letters*, 5(6), 709–714. <https://doi.org/10.1046/j.1461-0248.2002.00375.x>
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F. O. (2012). The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Research*, 41(D1), D590–D596. <https://doi.org/10.1093/nar/gks1219>
- Rohland, N., & Reich, D. (2012). Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Research*, 22(5), 939–946. <https://doi.org/10.1101/gr.128124.111>
- San Juan, P. A., Castro, I., & Dhami, M. K. (2021). Captivity reduces diversity and shifts composition of the Brown Kiwi microbiome. *Animal Microbiome*, 3(1), 1–8. <https://doi.org/10.1186/s42523-021-00109-0>
- San Juan, P. A., Hendershot, J. N., Daily, G. C., & Fukami, T. (2020). Land-use change has host-specific influences on avian gut microbiomes. *The ISME Journal*, 14(1), 318–321. <https://doi.org/10.1038/s41396-019-0535-4>
- Schmiedová, L., Tomášek, O., Pinkasová, H., Albrecht, T., & Kreisinger, J. (2022). Variation in diet composition and its relation to gut microbiota in a passerine bird. *Scientific Reports*, 12(1), 1–13. <https://doi.org/10.1038/s41598-022-07672-9>
- Skeen, H. R., Cooper, N. W., Hackett, S. J., Bates, J. M., & Marra, P. P. (2021). Repeated sampling of individuals reveals impact of tropical and temperate habitats on microbiota of a migratory bird. *Molecular Ecology*, 30(22), 5900–5916. <https://doi.org/10.1111/mec.16170>
- Sommer, F., & Bäckhed, F. (2013). The gut microbiota—Masters of host development and physiology. *Nature Reviews Microbiology*, 11(4), 227–238. <https://doi.org/10.1038/nrmicro2974>
- Song, S. J., Sanders, J. G., Delsuc, F., Metcalf, J., Amato, K., Taylor, M. W., Mazel, F., Lutz, H. L., Winker, K., Graves, G. R., Humphrey, G., Gilbert, J. A., Hackett, S. J., White, K. P., Skeen, H. R., Kurtis, S. M., Withrow, J., Braile, T., Miller, M., ... Knight, R. (2020). Comparative analyses of vertebrate gut microbiomes reveal convergence between birds and bats. *MBio*, 11(1), e02901-19. <https://doi.org/10.1128/mBio.02901-19>
- Suzuki, T. A. (2017). Links between natural variation in the microbiome and host fitness in wild mammals. *Integrative and Comparative Biology*, 57(4), 756–769. <https://doi.org/10.1093/icb/ixc104>
- Taberlet, P., Coissac, E., Pompanon, F., Brochmann, C., & Willerslev, E. (2012). Towards next-generation biodiversity assessment using DNA metabarcoding. *Molecular Ecology*, 21(8), 2045–2050. <https://doi.org/10.1111/j.1365-294X.2012.05470.x>
- Teysier, A., Matthysen, E., Hudin, N. S., De Neve, L., White, J., & Lens, L. (2020). Diet contributes to urban-induced alterations in gut microbiota: Experimental evidence from a wild passerine. *Proceedings of the Royal Society B*, 287(1920), 20192182. <https://doi.org/10.1098/rspb.2019.2182>
- Toews, D. P. L., Brelsford, A., Grossen, C., Milá, B., & Irwin, D. E. (2016). Genomic variation across the yellow-rumped warbler species complex. *The Auk: Ornithological Advances*, 133(4), 698–717. <https://doi.org/10.1642/AUK-16-61.1>
- Toews, D. P. L., Kramer, G. R., Jones, A. W., Brennan, C. L., Cloud, B. E., Andersen, D. E., Lovette, I. J., & Streby, H. (2020). Genomic identification of intergeneric hybrids in New World wood-warblers (Aves: Parulidae). *Biological Journal of the Linnean Society*, 131(1), 183–191. <https://doi.org/10.1093/biolinnean/blaa085>
- Toews, D. P. L., Streby, H. M., Burket, L., & Taylor, S. A. (2018). A wood-warbler produced through both interspecific and intergeneric hybridization. *Biology Letters*, 14(11), 20180557. <https://doi.org/10.1098/rsbl.2018.0557>
- Trevelline, B. K., Sosa, J., Hartup, B. K., & Kohl, K. D. (2020). A bird's-eye view of phyllosymbiosis: Weak signatures of phyllosymbiosis among all 15 species of cranes. *Proceedings of the Royal Society B*, 287(1923), 20192988. <https://doi.org/10.1098/rspb.2019.2988>
- Tung Ho, L. S., & Ané, C. (2014). A linear-time algorithm for Gaussian and non-Gaussian trait evolution models. *Systematic Biology*, 63(3), 397–408. <https://doi.org/10.1093/sysbio/syu005>
- Videvall, E., Song, S. J., Bensch, H. M., Strandh, M., Engelbrecht, A., Serfontein, N., Hellgren, O., Olivier, A., Cloete, S., Knight, R., & Cornwallis, C. K. (2019). Major shifts in gut microbiota during development and its relationship to growth in ostriches. *Molecular Ecology*, 28(10), 2653–2667. <https://doi.org/10.1111/mec.15087>
- Vo, A. T., & Jedlicka, J. A. (2014). Protocols for metagenomic DNA extraction and Illumina amplicon library preparation for faecal and

- swab samples. *Molecular Ecology Resources*, 14(6), 1183–1197. <https://doi.org/10.1111/1755-0998.12269>
- Volf, J., Polansky, O., Sekelova, Z., Velge, P., Schouler, C., Kaspers, B., & Rychlik, I. (2017). Gene expression in the chicken caecum is dependent on microbiota composition. *Veterinary Research*, 48(1), 85. <https://doi.org/10.1186/s13567-017-0493-7>
- White, P. S., & Densmore, L. D. (1992). Mitochondrial DNA isolation. In A. R. Hoelzel (Ed.), *Molecular genetic analysis of populations: A practical approach* (pp. 50–51). IRL Press at Oxford University Press.
- Xiao, K., Fan, Y., Zhang, Z., Shen, X., Li, X., Liang, X., Bi, R., Wu, Y., Zhai, J., Dai, J., Irwin, D. M., Chen, W., & Shen, Y. (2021). Covariation of the gut microbiome with diet in nonpasserine birds. *mSphere*, 6, e00308-21. <https://doi.org/10.1128/mSphere.00308-21>
- Youngblut, N. D., Reischer, G. H., Walters, W., Schuster, N., Walzer, C., Stalder, G., Ley, R. E., & Farnleitner, A. H. (2019). Host diet and evolutionary history explain different aspects of gut microbiome diversity among vertebrate clades. *Nature Communications*, 10(1), 2200. <https://doi.org/10.1038/s41467-019-10191-3>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Baiz, M. D., Benavides C., A., Miller, E. T., Wood, A. W., & Toews, D. P. L. (2023). Gut microbiome composition better reflects host phylogeny than diet diversity in breeding wood-warblers. *Molecular Ecology*, 32, 518–536. <https://doi.org/10.1111/mec.16762>